



# The Open Diabetes Journal

Content list available at: [www.benthamopen.com/TODIAJ/](http://www.benthamopen.com/TODIAJ/)

DOI: 10.2174/1876524601808010013, 2018, 8, 13-21



## CLINICAL TRIAL STUDY

### Type 2 Diabetes Study, Introduction and Perspective

Da-Yong Lu<sup>1,\*</sup>, Jin-Yu Che<sup>1</sup>, Nagendra Sastry Yarla<sup>2</sup>, Hong Zhu<sup>3</sup>, Ting-Ren Lu<sup>1</sup>, Bin Xu<sup>4</sup> and Swathi Putta<sup>5</sup>

<sup>1</sup>Shanghai University, Shanghai 200444, P. R. China

<sup>2</sup>Divisions of Biochemistry & Chemistry, City University of New York School of Medicine, 160 Convent Avenue, New York, NY 10031, USA

<sup>3</sup>College of Pharmaceutical Science, Zhejiang University, Hangzhou, P. R. China

<sup>4</sup>Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, P. R. China

<sup>5</sup>College of Pharmaceutical Science, Andhra University, Visakhapatnam, AP, India

Received: May 25, 2018

Revised: July 5, 2018

Accepted: July 16, 2018

#### Abstract:

##### Background:

The prevalence of Type 2 Diabetes Mellitus (T2DM) continues to rise globally. The T2DM prevalence is not only in developing countries, but also in developed countries now. Correspondingly, the therapeutics of T2DM calls for a change (higher efficiency) due to growing number of patients and increasing economic burdens globally. Entering into this millennium, both piecemeal pathways (idea driven) and exponential growth of human genomic study are developing quickly. Genotypic-phenotypic translation, modern diagnostics, pharmacology, drug developments, traditional Chinese medicine, personalized medicine and so on are promising disciplines for this change. The clinical anti-diabetic therapeutics, pathogenesis, drug development pipelines are especially highlighted.

##### Conclusion:

In summary, a general landscape and principle of T2DM is provided.

**Keywords:** Anti-diabetic therapy, Insulin-derivatives, Natural chemotherapeutic drugs, Diabetes-induced complication, Traditional Chinese Medicine, Diabetic pathology, Herbal medicine.


## 1. INTRODUCTION

The prevalence of Type 2 Diabetes Mellitus (T2DM) continues to rise globally. The T2DM prevalence is not only in developing countries, but also in developed countries now [1 - 4]. Correspondingly, the therapeutics of T2DM calls for a change (higher efficiency) due to growing number of patients and increasing economic burdens globally. Entering into this millennium, both piecemeal pathways (idea driven) and exponential growth of human genomic study are developing quickly. Genotypic-phenotypic translation, modern diagnostics, pharmacology, drug developments, traditional Chinese medicine, personalized medicine and so on are promising disciplines for this change. The clinical anti-diabetic therapeutics, pathogenesis, drug development pipelines are especially highlighted.

## 2. A COMMON BIOLOGICAL PROCESS FOR T2DM

In order to increase the efficiency of T2DM therapeutics, the basic knowledge of T2DM may be better understood.

\* Address correspondence to this author at the Shanghai University, Shanghai 200444, P. R. China, E-mail: [ludayong@shu.edu.cn](mailto:ludayong@shu.edu.cn)



**Fig. (1).** The most frequent pathway for T2DM induction.

In the long-course of T2DM induction and progress (Fig. 1), it is shown that a great number of risk-factors can totally play critical roles of disease progress.

Generally speaking, it was found that many other factors (chemical, biological or environmental influence) could induce the T2DM occurrence in animal model establishment [4 - 6] (Table 1). A line of chemical, genetic, cellular, physiological and pathological evidence, especially genetic and environmental aspects is tabulated in Table 1.

### 2.1. Importance of Early Diagnostics

Development of Type 2 Diabetes Mellitus (T2DM) involves a lengthy and complex process from disease onset into symptom/complication manifesting in susceptible human beings. As a result, the cascade characters must be noticed at the earliest in patients.

In the early stage, many preventive measures (regular exercises, non-smoking and so on) or diet control can slow down the pace of the ongoing stages. Some far-reaching medical education systems for diabetes should be introduced to human beings in order to get medical intervention as early as possible [1]. Due to the asymptomatic characteristics in early stages of metabolic syndromes, early technical diagnostics is inevitable yet imperfect now. People should be

aware that high calorie food consumption is not the only factor to induce T2DM, some other unfavorable risk factors, such as habitually heavy drinking, somatic mutations, toxic compounds or sleep problems *etc.* [4 - 6]. can also trigger the incidence of T2DM. Thus, medical check-up for blood glucose concentration and levels should be carried out for people more than 40 years old every year.

**Table 1. Possible categories for T2DM initiations and causality.**

Etiology	Possible Pathological Pathways	References
Genetic/epigenetic	DNA methylation Copy-number change Gene transcription Loss-of-function mutation	[3 - 8]
Molecular	Growth factor excretion Leptin Insulin Adipose-hypoxia Adipocyte-respiration Pituitary hormonal synthesis and secretion	[7 - 9]
Chemical	Toxic compounds Other drugs	[10]
Physiology	Old age Liver dysfunction Inflammation Obesity Insulin-resistance Sleep problems Energy homeostasis Hypothalamic circuit Immune-metabolic pathways	[7 - 9, 11 - 18]
Food	More energy foods Palatable food (processed foods) Free-accessed foods	[11 - 13]
Life-style Behavior	Sedentary occupation Exercise deficiency Long-term TV watching Alcoholic Substance-addictive	[4, 5, 11 - 16]

**2.2. Therapeutic Ideas in General**

Regular patho-therapeutic ideas and current drug targets are enlisted as following pathways;

1. What is the therapeutic outcome and differences between patients with T2DM of various origins?
2. What are genetic-molecular relationships between blood glucose increase and disease complications?
3. May a known variety of T2DM origins relevant to different individuals?
4. Can we successfully predict what kinds of diabetic-induced complications will emerge under specific conditions of patients?

**3. GENETIC AND MOLECULAR ASPECT OF T2DM**

In order to update therapeutics, questions of whether genetic, molecular, physiological, behavioral and environmental aspects of disease-association pathways might be answered and observed;

1. By taking account individual variability in genetics, environmental and life-style, might some effective medications be separately validated?
2. What is the core of different anti-diabetic therapeutics?
3. Can curable therapeutics be widely achieved by patients with T2DM?


Without fully understanding the patho-therapeutic relation, we will not be able to retard diabetic-induced complications as early as possible. These avenues are inevitable for type 2 diabetes study. Several topics are described

in this respect.

### 3.1. Human Genomic Study

T2DM is generated from genetic factors interaction with environmental influence. Personalized anti-Diabetic Therapy (PDT) may be a modern possibility that enables us to offer better therapeutics by integration of different technology and drugs. Pharmacogenetics (PG) [19], the widest utility of PDT has a great potentiality for drug selecting and dosing. To accomplish PG for T2DM treatments, human genetic/pathologic information must be known before therapeutics. With the advent and maturation of genomic sequencing technology [20], this kind of human genomic study might become practical to all patients and get information exponentially [20, 21]. But, who has the right to use human genomic information without regulation [22]. It is an ethical question that needs long-term debates between patients and healthcare providers. Currently, the “loss-of-function” by genetic mutation, such as leptin, insulin and their receptors or down-stream molecules is widely observed [21].

At this stage of technical capability, PG study for diabetes widely encompasses drug dosing (pharmacodynamics or pharmacokinetics issue) by genotypes of human metabolic enzymes. Due to the unpredictable nature of overall disease-related molecular alterations and metabolic abnormality, PG for various diabetic profiling is the important information different from metabolic enzymes (Fig. 2). Fig. (2) represents the outlook of PDT applications and promotion in the clinic. This type of PDT may be utilized in anti-diabetic treatments in the future [19 - 21].



**Fig. (2).** Proposed diagram of personalized anti-diabetic therapy.

### 3.2. Molecular and Physiologic Pathways and Influences

Molecular and physiologic pathways and influences for T2DM are exemplified as the following parts;

Neural activity and circuits that control the appetite of patients is now located onto brain hypothalamic-pituitary

area [17]. These kinds of brain activities are very important for human obesity and T2DM.

Abnormal metabolic conditions in human adipose tissues; their dysfunctional characteristics are based on adipose tissues, livers and other human organs [18].

Hormonal synthesis, secretion and resistance, such as insulin, leptin and others play key roles for patients' obesity and diabetes [4 - 18]. Their synthesis, secretions and functional resistance are useful for individualized treatments of T2DM.

Energy homeostasis; a lot of patients with obesity and T2DM requires food control and heightened exercise activities [15, 16]. Human energy balance will decide human body weight gain or loss and further affect blood glucose levels. This energy issue is crucial for T2DM progress and managements [21].

Apart from these disease-induced pathways, other genetic and molecular issues may also involve [1 - 4]. With the growth of T2DM therapeutic study, higher efficiency of disease controls may be expected.

#### 4. PHARMACEUTICAL AND PHARMACOLOGIC STUDIES

##### 4.1. Herbal Drugs and Traditional Chinese Medicine (TCM)

A great variety of herbs (>500 species) have been reported to be associated with T2DM [10]. A lot of articles reported their functionality and therapeutic activity in experimental settings. Nonetheless, only a few of these herbal drugs have been commercialized and widely utilized for T2DM prevention and treatments. Likewise, herbal therapy is one of the major avenues of *diabetes mellitus* therapeutic study and clinical development in the future [1, 23].

Most therapeutic data from anti-diabetic herbs are commonly based on multitude biochemical pathways, such as antioxidant activity, insulin synthesis, secretion and sensitivity, glucose uptake assistance, reduced glucose production from hepatocytes, anti-inflammatory activity and so on [22]. However, except a small number of countries, such as China, India and Germany, herbal drugs are not allowed for therapeutic purpose worldwide because of the requisite of high-quality and complexity of drug prescription in the clinic. From pharmacological perspective, the therapeutic study for herbal medicines should aim at specific drug targets and whole body regulation in clinical trials. Many animal diabetic models might not be the best vehicle for direct translations in the clinic.

To better utility of herbal drugs or recipes, some principle of TCM for diabetes might be properly established in the future [1]. Combinations of anti-diabetic therapy of modern medicine (chemical or biological therapy) and traditional (TCM) might be useful in the clinic.

##### 4.2. Pharmaceutical Study

Insulin-based therapy needs to improve for clinical feasibility. It commonly needs syringes and human-body injections, which lead to treatment inconvenience. Overall, some small-molecular chemicals having the insulin-configurations or insulin receptors binding activity might be potential oral anti-diabetic drugs for improving insulin-based therapy in the clinic [13, 14].

More recently, some capsules containing biological form of insulin were invented for patients needing insulin therapy [1]. This pharmaceutical innovation can provide greater feasibility for diabetic treatments in the future.

##### 4.3. Therapeutic Applications Nowadays

Approximately ten drugs have been widely used for T2DM worldwide. Table 2 outlines parts of these therapeutic drugs and potential applications in the clinic. Despite wide-ranges of drug developments, clinical T2DM treatments are commonly not curable [1, 23]. To most of these therapeutic options, drug treatments will be life-long for high-proportion of T2DM patients. After all, we need to know more about patho-therapeutic relations for curable therapy for patients with T2DM in the future.

**Table 2. The recommended therapeutic options in clinical diabetic treatments.**

Drug and therapy	Mechanisms of action	Adverse events
Insulin and its derivatives	Glucose metabolisms	Not significance
Sulphonylurea	Stimulate insulin secretions	Gastrointestinal (20-30%) Infections (20%)
Biguanide (Metformin)	Decrease amount of sugar productions by liver	Metabolisms

(Table 2) contd....

Drug and therapy	Mechanisms of action	Adverse events
Acarbose Voglibose	$\alpha$ -glucosidase inhibitors	Gastrointestinal (20-30%)
Pioglitazone	Receptor agonists	Heart failure (1-5%) Bladder carcinoma (1-2%)
Sitagliptin Vidagliotin Saxagliptin	Dipeptidyl peptidase IV inhibitors	Injection
GLP-1 analogues	Nature peptic C	Injection
Gene therapy	Insulin-formation	Under-investigation
Pancreatic transplantation	Recovery of pancreatic function	Difficult to implement
Exercise	Mimics insulin actions	Not significance
Curcumin Resveratrol	Plant chemicals Wider biochemical pathways	Low
Other melbine	Variable	Variability
Smoke cessation	Cardiovascular risks	No
Aspirin	Cardiovascular risks	Blood coagulation impair
Vaccine	Infections	Not obvious
Gastric bariatric surgery	Obesity	Expensive
Orlistat	Obesity (pancreatic lipase)	Only drug for obesity
Propolis	Disease complication	No
Natto	Vascular softness	No
Herbal drug (>500)	Variable	Variability
$\alpha$ -thioctic acids (Lipoic acid)	Disease complications	Not obvious
New anti-diabetic drugs	SGLT-2 inhibition DPP4 antagonists	Under investigations

[1, 24].

Apart from herbal drugs, some insect products (Propolis) [25, 26] and fermented soybean (Natto) [4, 5] were also applicable in China and Japan for T2DM-related symptom control.

#### 4.4. Therapeutic Combination in Clinical Trials

In the past, life-style adjustment is an effective way for the reductions of blood glucose and complication events for patient's notification [27 - 29]. As usual, combination of life-style adjustment and metformin is widely recommended by both American Diabetes Association (ADA) and European Association for Study of Diabetes (EAST). Table 2 shows a great number of therapeutic options chosen for anti-diabetic treatments. Certainly, we commonly use more than one therapeutic option in clinical trials. Generally, drug combination is more welcome for refractory chronic disease, like cancer and HIV/AIDS treatments comparing with mono-therapy [30 - 34]. It is a new concept of whether a systematic approach could be helpful from therapeutic combination.

### 5. ROAD MAP IN THE FUTURE

#### 5.1. Pathological Issues

Expanding pathological knowledge will be introduced to broader ranges of people for noticing and treating patients on disease onset. Genome Wide Association Study (GWAS) between patients and normal people may be breakthroughs for T2DM pathogenesis and treatments - the exponentially explosion of human genomic data.

#### 5.2. Therapeutic Issues

Genetic or molecular study of disease progression, drug pharmacology, mechanisms of action for drug activity/toxicity. Establish personalized anti-diabetic therapy in the clinic. Develop higher therapeutic-index drugs, such as natural chemotherapeutic anti-diabetic drugs [35, 36]. Pharmaceutically improves insulin-based treatments from needle to oral. Budget control in drug developments and cost-effective in clinical treatment selections will be more optimized [37].

#### 5.3. Regulatory Issue

Global cooperation is inevitable, especially invitation of mathematical majored students and scholar in this medical

campaign [38 - 40]. Mathematical solution of disease risks and occurrence is given in Equation 1 [41].

$$\mathbf{T} (P_1, \dots, P_n) = \boldsymbol{\theta} (P_n) \quad (1)$$

where, T: incidence and risks of T2DM in random human population

## CONCLUSION

Despite a lot of discoveries, current therapeutic agents (such as insulin and other drugs) are economic burden to patients due to long-term of drug utility (generally >400 USD per year). Rapid development of higher efficiency anti-diabetic drugs, especially for diabetic-induced complications, is a high priority and of great medical significance [42 - 46]. A step closer should be a marvelous thing.

Given too much molecular components and pathways can be involved in T2DM origins, growing volumes of financial supports must be allocated in both developing countries and developed countries. To sum up, win-win situation between developed countries and developed countries can decide how long we can go through in this medical campaign. After global cooperation, both extremes of human populations can be provided with high quality of anti-diabetic medications. In the future, new perspectives and scientific investigations will be introduced for changing the landscape of current anti-diabetic therapeutics if possible. A lot of therapeutic options need to be better served for complicated situations of diabetes in the clinic.

## ETHICAL APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

## HUMAN AND ANIMAL RIGHTS

No Animals/Humans were used for studies that are bases of this research.

## CONSENT FOR PUBLICATION

Not applicable.

## CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

## ACKNOWLEDGEMENTS

This work was funded by Shanghai Science and Technology Foundation of High Educations 97A49.

## REFERENCES

- [1] Lu DY, Che JY, Yarla NS, *et al.* Types 2 diabetes prevention, treatments and new drug developments. *Clin Immunol Endocr Metab Drugs* 2018; 5.
- [2] Kaul K, Tarr JA, Ahmad SI, Kohner EM, Chibber R. Introduction to diabetes Mellitus. *Diabetes: An Old Disease, A New Insights*. In: Ahmad S, Ed. Springer Science, US 2013; pp. 1-11.
- [3] Zimmet PZ, Magliano DJ, Herman WH, Shaw JE. Diabetes: A 21<sup>st</sup> century challenge. *Lancet Diabetes Endocrinol* 2014; 2(1): 56-64. [[http://dx.doi.org/10.1016/S2213-8587\(13\)70112-8](http://dx.doi.org/10.1016/S2213-8587(13)70112-8)] [PMID: 24622669]
- [4] Ahmad S, Ed. *An Old Disease, A New Insights*. Springer Science, US 2013.
- [5] Lu DY, Che JY, Yarla NS, *et al.* Diabetes prevention and treatments, a specific topic for modern medicines. *J Metab Syndr* 2017; 8(3): 231.
- [6] Lu DY, Che JY, Yarla NS, *et al.* Type 2 diabetes, medical knowledge and pharmaceutical innovations. *J Diabetol* 2017; 1(1): 1-3.
- [7] Yang W, Lu J, Weng J, *et al.* Prevalence of diabetes among men and women in China. *N Engl J Med* 2010; 362(12): 1090-101. [<http://dx.doi.org/10.1056/NEJMoa0908292>] [PMID: 20335585]
- [8] Fuchs S, Henschke C, Blümel M, Busse R. Disease management programs for type 2 diabetes in Germany: A systematic literature review evaluating effectiveness. *Dtsch Arztebl Int* 2014; 111(26): 453-63. [PMID: 25019922]
- [9] Putta S, Peluso I, Yarla NS, *et al.* Diabetes mellitus and male aging, pharmacotherapeutics and clinical implications. *Curr Pharm Des* 2017; 23(41): 6321-46. [<http://dx.doi.org/10.2174/1381612823666170519151801>] [PMID: 28741457]

- [10] Singh A, Srivastav R, Randey AK. Protective role of Terminalia Chebula in streptozotocin-induced diabetic mice for wound healing activity. *Br J Med Med Res* 2017; 22(2): 1-8.  
[<http://dx.doi.org/10.9734/BJMMR/2017/33137>]
- [11] Nannapaneni S, Ramar K, Surani S. Effect of obstructive sleep apnea on type 2 diabetes mellitus: A comprehensive literature review. *World J Diabetes* 2013; 4(6): 238-44.  
[<http://dx.doi.org/10.4239/wjd.v4.i6.238>] [PMID: 24379913]
- [12] Freire MDM, Alves C. Therapeutic Chinese exercises (Qigong) in the treatment of type 2 diabetes mellitus: A systematic review. *Diabetes Metab Syndr* 2013; 7(1): 56-9.  
[<http://dx.doi.org/10.1016/j.dsx.2013.02.009>] [PMID: 23517799]
- [13] Lu DY, Che JY, Wu HY, Lu TR. The pathogenesis and treatments of diabetes, questions and answers. *Cell Dev Biol* 2014; 3(3): e126.
- [14] Lu DY, Che JY, Wu HY, Lu TR. The pathogenesis and treatments of diabetes, a new insight. *Adv Tech Biol Med* 2014; 2: e102.
- [15] Lu DY, Che JY, Wu HY, *et al.* An overview of obesity. *Metabolomics* 2018; 8(2): 200.
- [16] Lu DY, Che JY, Wu HY, *et al.* Obesity, risks and managements. *Metabolomics* 2018; 8(1): e156.
- [17] Steculorum SM, Paeger L, Bremser S, *et al.* Hypothalamic UDP increases in obesity and promotes feeding *via* P2Y6-dependent activation of AgRP neurons. *Cell* 2015; 162(6): 1404-17.  
[<http://dx.doi.org/10.1016/j.cell.2015.08.032>] [PMID: 26359991]
- [18] Lee YS, Kim JW, Osborne O, *et al.* Increased adipocyte O<sub>2</sub> consumption triggers HIF-1 $\alpha$ , causing inflammation and insulin resistance in obesity. *Cell* 2014; 157(6): 1339-52.  
[<http://dx.doi.org/10.1016/j.cell.2014.05.012>] [PMID: 24906151]
- [19] Corrêa-Giannella ML, Machado UF. SLC2A4 gene: A promising target for pharmacogenomics of insulin resistance. *Pharmacogenomics* 2013; 14(8): 847-50.  
[<http://dx.doi.org/10.2217/pgs.13.45>] [PMID: 23746177]
- [20] Lander ES. Initial impact of the sequencing of the human genome. *Nature* 2011; 470(7333): 187-97.  
[<http://dx.doi.org/10.1038/nature09792>] [PMID: 21307931]
- [21] van der Klaauw AA, Farooqi IS. The hunger genes: Pathways to obesity. *Cell* 2015; 161(1): 119-32.  
[<http://dx.doi.org/10.1016/j.cell.2015.03.008>] [PMID: 25815990]
- [22] Rahimzadeh V, Bartlett G. Policies and practices of data-intensive primary care in the precision-medicine era. *Intern Med Rev* 2017; 3(9): 1-14.  
[<http://dx.doi.org/10.18103/imr.v3i9.558>]
- [23] Li GQ, Kam A, Wong KH, *et al.* Herbal medicines for the management of diabetes. An old disease, A new insights. In: Ahmad S, Ed. Springer Science, US 2013; pp. 396-413.
- [24] Imam K. Management and treatment of diabetes mellitus. An old disease, A new insights. In: Ahmad S, Ed. Springer Science, US 2013; pp. 356-80.
- [25] Lu DY, Che JY. Rethink of diabetes treatment and drug development. *Cell Dev Biol* 2014; 3(2): e125.
- [26] Wagh VD. Propolis: A wonder bees product and its pharmacological potentials. *Adv Pharmacol Sci* 2013; 2013: 308249.  
[<http://dx.doi.org/10.1155/2013/308249>] [PMID: 24382957]
- [27] Thepwoonga I, Kirby C, Schattner P, Shaw J, Piterman L. Type 2 diabetes continuing medical education for general practitioners: What works? A systematic review. *Diabet Med* 2014; 31(12): 1488-97.  
[<http://dx.doi.org/10.1111/dme.12552>] [PMID: 25047877]
- [28] Grimaccia F, Kanavos P. Cost, outcomes, treatment pathways and challenges for diabetes care in Italy. *Global Health* 2014; 10(1): 58.  
[<http://dx.doi.org/10.1186/1744-8603-10-58>] [PMID: 25024101]
- [29] Wens J, Vermeire E, Hearnshaw H, Lindenmeyer A, Biot Y, Van Royen P. Educational interventions aiming at improving adherence to treatment recommendations in type 2 diabetes: A sub-analysis of a systematic review of randomised controlled trials. *Diabetes Res Clin Pract* 2008; 79(3): 377-88.  
[<http://dx.doi.org/10.1016/j.diabres.2007.06.006>] [PMID: 17643546]
- [30] Lu DY, Lu TR, Cao S. Drug combinations in cancer treatment. *Clin Exp Pharmacol* 2013; 3: 134.
- [31] Lu DY, Chen EH, Lu TR, Ding J. Anticancer drug combinations, studies from different pathways. *Cell Dev Biol* 2015; 4(5): 166.
- [32] Lu DY, Wu HY, Yarla NS, Xu B, Ding J, Lu TR. HAART in HIV/AIDS treatments, future trends. *Infect Disord Drug Targets* 2018; 18(1): 15-22.  
[<http://dx.doi.org/10.2174/1871526517666170505122800>] [PMID: 28474549]
- [33] Lu DY, Chen EH, Wu HY, Lu TR, Xu B, Ding J. Anticancer drug combination, how far we can go through? *Anticancer Agents Med Chem* 2017; 17(1): 21-8.  
[PMID: 27039923]
- [34] Lu DY, Lu TR, Yarla NS, *et al.* Drug combination in clinical cancer treatment. *Rev Recent Clin Trials* 2017; 12(3): 202-11.  
[<http://dx.doi.org/10.2174/1574887112666170803145955>] [PMID: 28782482]



- [35] Lu DY, Lu TR, Lu Y, Sastry N, Wu HY. Discover natural chemical drugs in modern medicines. *Metabolomics* 2016; 6(3): 181.
- [36] Alam F, Islam MA, Kamal MA, Gan SH. Updates on managing type 2 diabetes mellitus with natural products. Towards antidiabetic drug developments. *Curr Med Chem* 2016; 23(29): 1-37. [http://dx.doi.org/10.2174/0929867323666160813222436] [PMID: 27528060]
- [37] Asche C, LaFleur J, Conner C. A review of diabetes treatment adherence and the association with clinical and economic outcomes. *Clin Ther* 2011; 33(1): 74-109. [http://dx.doi.org/10.1016/j.clinthera.2011.01.019] [PMID: 21397776]
- [38] Lu DY, Lu TR, Lu Y, Wu HY, Yarla NS. The acquisition of mathematical language in biomedical articles. *J Cell Developmental Biol* 2017; 1(1): 8.
- [39] Lu DY, Lu TR. Mathematics or physics-majored students on the biomedical fields, insiders or outsiders? *Metabolomics* 2015; 5(4): e142.
- [40] Lu DY, Wu HY, Lu TR, Che JY, Lu Y. Updating biomedical studies by recruiting more mathematics or physics-majored talents. *Metabolomics* 2016; 6(2): e148.
- [41] Gentle JE. *Elements of Computational Statistics*. Germany: Springer Science 2002.
- [42] Turpin SM, Hoy AJ, Brown RD, *et al*. Adipose triacylglycerol lipase is a major regulator of hepatic lipid metabolism but not insulin sensitivity in mice. *Diabetologia* 2011; 54(1): 146-56. [http://dx.doi.org/10.1007/s00125-010-1895-5] [PMID: 20842343]
- [43] Putta S, Yarla NS, Peluso I, *et al*. Anthocyanins: Possible role as multitarget therapeutic agents for prevention and therapy of chronic diseases. *Curr Pharm Des* 2017; 23(30): 4475-83.
- [44] Davis TME, Ting R, Best JD, *et al*. Effects of fenofibrate on renal function in patients with type 2 diabetes mellitus: The fenofibrate intervention and event lowering in diabetes (FIELD) study. *Diabetologia* 2011; 54(2): 280-90. [http://dx.doi.org/10.1007/s00125-010-1951-1] [PMID: 21052978]
- [45] Metabolex Inc. Use of (-) (3-trihalomethylphenoxy)(4-halophenyl) acetic acid derivatives for treatment of type 2 diabetes. US8329749 (2012).
- [46] Metabolex Inc. N-linked heterocyclic receptor agonists for the treatment of diabetes and metabolic disorders. US20120322804A1 (2012).

© 2018 Lu *et al*.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.